



GOALS OF CARE DESIGNATION (GCD) ORDER

DATE	TIME
_____	_____
(YYYY) (MM) (DD)	(hh:mm)

Goals of Care Designation Order
 To order a Goals of Care Designation for this patient, check the appropriate Goals of Care Designation below and write your initials on the line below it.
(See reverse side for details)

Check ▶ <input type="checkbox"/> R1	<input type="checkbox"/> R2	<input type="checkbox"/> R3	<input type="checkbox"/> M1	<input type="checkbox"/> M2	<input type="checkbox"/> C1	<input type="checkbox"/> C2
Initials ▶ _____	_____	_____	_____	_____	_____	_____

Check (✓) here if this GCD Order is an interim Order awaiting the outcome of a Dispute Resolution Process.
 Document further details on the ACD/GCD Tracking Record as well.

Specify here if there are specific clarifications to this GCD Order. Document these clarifications on the ACP/GCD Tracking Record as well.

Patient's location of care where this GCD Order was ordered *(Home; or clinic or facility name)*

Indicate which of the following apply regarding involvement of the Patient or alternate decision-maker (ADM)

This GCD has been ordered after relevant conversation with the patient.

This GCD has been ordered after relevant conversation with the alternate decision-maker (ADM), or others.
(Names of formally appointed or informal ADM's should be noted on the ACP/GCD Tracking Record)

This GCD is an interim GCD Order prior to conversation with patient or ADM.

History/Current Status of GCD Order

Indicate one of the following

This is the first GCD Order I am aware of for this patient.

This GCD Order is a revision from the most recent prior GCD *(see ACP/GCD Tracking Record for details of previous GCD Order)*

This GCD Order is unchanged from the most recent prior GCD.

Name of Physician / Designated Most Responsible Health Practitioner who has ordered this GCD	Discipline

Signature	Date (yyyy/mm/dd)

Goals of Care Designation – Guide for Clinicians

The Goals of Care Designation provides direction regarding specific health interventions, transfer decisions, locations of care, and limitations on interventions for a Patient as established after consultation between the Most Responsible Health Professional and Patient.

The Windsor Regional Hospital Goals of Care Designations replace the levels of care (“code levels”) identified in the former Adult Resuscitation. The Goals of Care Designations are INDEPENDENT of the Patient’s current location. The Designation Order follows a Patient in order to guide the “receiving” health care providers if new conditions occur. However, flexibility to make appropriate and altered clinical decisions given new information or new conditions is always retained.

Transfer of a Patient from long term or supportive living care to Acute Care is a consideration in determining a Designation. The goal of such a transfer is aimed at cure or control of the medical condition. A decision not to transfer under such circumstances implies that, if the Patient’s condition worsens despite treatments in the Patient’s current environment and becomes irreversible, a mode of care focusing on comfort and symptom control is adopted and a natural death occurs. Transfer may still occur for such Patients if the goals is to investigate or treat symptoms, and if efforts aimed at this is best undertaken at another location.

<p>R: Medical Care and Interventions, Including Resuscitation if required followed by Intensive Care Unit Admission. Focus of Care and interventions are for cure or control of the Patient’s condition. The Patient would desire and is expected to benefit from attempted resuscitation and ICU care if required</p> <p>R1: Patient is expected to benefit from and is accepting of any appropriate Investigations/Interventions that can be offered including attempted resuscitation and ICU care</p> <ul style="list-style-type: none"> • Resuscitation: is undertaken for acute deterioration, and may include intubation and chest compression • Life Support Interventions: are usually undertaken • Life Sustaining Measures: are used when appropriate • Major Surgery: is considered when appropriate • Transfer: is considered for diagnosis and treatment, if required <p>R2: Patient is expected to benefit from and is accepting of any appropriate investigations/interventions that can be offered including attempted resuscitation, intubation and ICU care, but excluding chest compression</p> <ul style="list-style-type: none"> • Resuscitation: is undertaken for acute deterioration, but chest compression should not be performed. • Life Support Interventions: may be offered without chest compression • Life Sustaining Measures: are used when appropriate • Major Surgery: is considered when appropriate • Transfer: is considered for diagnosis and treatment, if required <p>R3: Patient is expected to benefit from and is accepting of any appropriate investigations/interventions that can be offered including attempted resuscitation and ICU care, but excluding intubation and chest compression</p> <ul style="list-style-type: none"> • Resuscitation: is undertaken for acute deterioration but intubation and chest compression should not be performed • Life Support Interventions: may be offered without intubation and without chest compression • Life Sustaining Measures: are used when appropriate • Major Surgery: is considered when appropriate • Transfer: is considered for diagnosis and treatment, if required 	<p>M: Medical Care and Interventions, Excluding Resuscitation. Focus of Care and interventions are for cure or control of the Patient’s condition. The Patient either chooses to not receive or would not be expected to benefit from attempted resuscitation followed by life-sustaining care in an ICU. In Pediatrics, ICU can be considered if that location is deemed the best location for delivery of specific short-term <u>symptom</u>-directed care.</p> <p>M1: All clinically appropriate medical and surgical interventions directed at cure and control of condition(s) are considered, excluding the option of attempted life-saving resuscitation followed by ICU care. See above, regarding Pediatrics and ICU.</p> <ul style="list-style-type: none"> • Resuscitation: is not undertaken for cardio respiratory arrest. • Life Support Interventions: should not be initiated, or should be discontinued after discussion with the Patient. • Life Sustaining Measures: are used when appropriate • Transfer: to another location of care is considered if that location provides more appropriate circumstances for diagnosis and treatment. • Major Surgery: is considered when appropriate. Resuscitation during surgery or in the recovery room can be considered, including short term physiologic and mechanical support in an ICU, in order to return the Patient to prior level of function. The possibility of intra-operative death or life-threatening deterioration should be discussed with the Patient in advance of the proposed surgery and general decision-making guidance agreed upon and documented. <p>M2: All clinically appropriate interventions that can be offered in the current <u>non-hospital</u> location of care are considered. If a patient does not respond to available treatments in this location of care, discussion should ensue to change the focus to comfort care. Life-saving resuscitation is not undertaken except in unusual circumstances (see below in Major Surgery) See above, regarding Pediatrics and ICU.</p> <ul style="list-style-type: none"> • Resuscitation: is not undertaken for cardio respiratory arrest. • Life Support Interventions: should not be initiated, or should be discontinued after discussion with the Patient. • Life Sustaining Measures are: used when appropriate • Transfer: is not usually undertaken but can be contemplated if symptom management or diagnostic efforts aimed at understanding symptoms can be best undertaken at that other location. • Major Surgery: can be considered, in order to prevent suffering from an unexpected trauma or illness. Resuscitation during surgery or in the recovery room can be considered, including short term physiologic and mechanical support in an ICU, in order to return the Patient to prior level of function. The possibility of intra-operative death or life-threatening deterioration should be discussed with the Patient in advance of the proposed surgery and general decision-making guidance agreed upon and noted as special circumstances on the GCD Order Form and Tracking Record. 	<p>C: Medical Care Interventions, Focused on Comfort. Focus of Care and interventions are for the active palliative treatment of the Patient who has a terminal illness, and support for those close to them. This includes medical care for symptom control and psychosocial and spiritual support in advance of death. Care can be provided in any location best suited for these aims, including an ICU, a Hospice or any location that is the most appropriate for symptom-based care for this particular Patient.</p> <p>C1: All care is directed at maximal symptom control and maintenance of function without cure or control of an underlying condition that is expected to cause eventual death. Treatment of intercurrent illnesses can be contemplated only after careful discussion with the Patient about specific short-term goals.</p> <ul style="list-style-type: none"> • Resuscitation: is not undertaken. • Life Support Interventions: should not be initiated, or should be discontinued after discussion with the Patient. • Life Sustaining Measures: are used only for goal directed symptom management. • Major Surgery: is not usually undertaken but can be contemplated for procedures aimed at symptom relief. Resuscitation during surgery or in the recovery room can be considered, including short term physiologic and mechanical support in an ICU, in order to return the Patient to prior level of function, but this would be a rare circumstance. The possibility of intra-operative death or life-threatening deterioration should be discussed with the Patient in advance of the proposed surgery and general decision-making guidance agreed upon and documented. • Transfer: to any appropriate location of care can be considered at any time, to better understand or control symptoms. <p>C2: All care is directed at preparation for imminent death [usually within hours or days] with maximal efforts directed at symptom control.</p> <ul style="list-style-type: none"> • Resuscitation: is not undertaken. • Life Support Interventions: should not be initiated, or should be discontinued after discussion with the Patient. • Life Sustaining Measures: should be discontinued unless required for symptom management. • Major Surgery: is not appropriate. • Transfer: is usually not undertaken but may be considered if required.
---	---	---